BETWEEN THE ROWS®

Performing Yield Estimates

Wyffels.com August 12, 2025

In late summer and early fall, many corn growers evaluate fields and make yield estimates either for storage or marketing purposes. Estimating yield can be difficult since weather conditions can result in highly variable yield differences within a field. To improve accuracy of yield estimates, multiple samples need to be taken throughout a field.

Estimate on a whole field basis

When conducting whole field yield estimates, the Yield Component Method is commonly used and is calculated by:

 $\frac{\text{# of harvestable ears per ac X \# kernels per ear}}{\text{# of kernels per bushel}} = \text{Yield (bu/ac)}$

This yield estimate is best done at least 3 weeks after pollination when kernels are in the milk stage (R3) and beyond.

Kernels per bushel

The number of kernels required to make a bushel can be highly variable among hybrids and growing conditions. Typically, 90,000 kernels per bushel is used as an average, though hybrid differences are known to vary by over 30%.

Each year Wyffels research calculates the number of kernels per bushel (kpb) for each genetic family (see table on page 2). This factor will vary based on the growing conditions each year. For instance, kernel size tends to be smaller in lower yield environments which increases the kpb factor. Conversely, kernel size is typically larger in higher yield environments due to ideal conditions during grainfill, thus the kpb factor is smaller.

Although the exact number of kernels per bushel for each hybrid will vary each year based on crop conditions, the relative differences between hybrids will remain fairly consistent. So, even in more variable conditions the listed kernel per bushel factors will serve as a great tool to see differences between hybrids when performing yield estimates.

Yield estimate protocol

Use the following protocol to ensure more accurate estimates.

Ear counts

- Measure 1/1000th of an acre in one row (17'5" for 30inch rows) and count the number of harvestable ears present. Do not rely on planting population as not all seeds likely emerged and every plant might not have an ear.
- Take several ear counts across the field to represent all the conditions present. Don't just hit the garden spots. In fields impacted by heat and drought or that have reduced stands, ten or more samples will be needed to ensure reasonable representation of the entire field.

Kernel counts

- Collect 3 to 5 ears in each harvestable ear count.
- Collect every 7th ear to get an unbiased, representative sample.
- Count and multiply the number of kernel rows around by the number of kernels in a row for each ear. For example: 16 kernel rows around x 34 kernels per row = 544 kernels/ear.
- Average the kernel counts across the ear samples.

Calculate yield

- Plug in the harvestable ears per acre and kernels per ear into the above formula
- Plug in the kernel per bushel factor for listed hybrids. If the hybrid is not listed, use 90,000 kpb for average sized kernels, 110,000 kpb if kernels are small, and 70,000 kpb for large kernels.

Wyffels Yield Calculator

In order to make yield estimates easier and more efficient, Wyffels has developed an online yield calculator. The Wyffels Yield Calculator can be found on the Wyffels.com website under the Agronomy tab. This calculator has been pre-programmed with all of the specific hybrid factors for hybrids in the Wyffels lineup. There is also a yield environment option to account for kernel size variability. To use the calculator, **click this link.**



YIELD FACTOR

Important note on hybrid kernel factors: These factors represent an average observation for each hybrid, and will be most accurate in yield environments of 210-250 bu/A.

Genetic Famil	y Factor	Genetic Fam	ily Factor
W1306	84	W5129	80
W1356	80	W5406	72
W1546	80	W5518	73
W1548	79	W5684	71
W1599	78	W5778	74
W1676	80	W6110	73
W1758	75	W6215	71
W1826	85	W6408	75
W1869	78	W6479	80
W1996	76	W6505	66
W1988	84	W6630	73
W2016	80	W6826	72
W2236	76	W6886	71
W2386	75	W6935	66
W2446	78	W7048	66
W2506	76	W7170	70
W2557	84	W7208	77
W2595	78	W7416	66
W2648	74	W7424	70
W2656	78	W7456	65
W2629	76	W7485	66
W2674	73	W7499	69
W3095	69	W7510	66
W3286	74	W7536DG	76
W3309	71	W7696	72
W3576	68	W7726	75
W3579	69	W7759	73
W3654	73	W7878	66
W4025	75	W7888	75
W4196	70	W7945	70
W4246	75	W7956	64
W4358	72	W8007	73
W4439	69	W8086	66
W4487	72	W8108	74
W4514	73	W8306	67
W4796	74	W8936DG	68
W5019	72	W9218	70
W5086	72		

Summary

No matter how detailed you are with this process, remember that you are performing an estimate. To improve accuracy, increase your sample number throughout the field. Using the kernels per bushel factor for each hybrid will help fine tune estimates for your Wyffels products. And using the online Wyffels Yield Calculator will help reduce mathematical error.

From the desk of



Dr. Brent Tharp Technical Product Manager

309.737.6434 cell btharp@wyffels.com

